

11155 Be Efficient

Consider an integer sequence consisting of N elements, where:

$$\begin{aligned}
 X_0 &= A \\
 X_i &= ((X_{i-1} * B + C) \% M) + 1 \quad \text{for } i = 1 \text{ to } N - 1
 \end{aligned}$$

You will be given the values of A , B , C , M and N . Find out the number of consecutive subsequences whose sum is a multiple of M .

Consider an example where $A = 2$, $B = 1$, $C = 2$, $M = 4$ and $N = 4$.

So, $X_0 = 2$, $X_1 = 1$, $X_2 = 4$ and $X_3 = 3$.

The consecutive subsequences are $\{2\}$, $\{2\ 1\}$, $\{2\ 1\ 4\}$, $\{2\ 1\ 4\ 3\}$, $\{1\}$, $\{1\ 4\}$, $\{1\ 4\ 3\}$, $\{4\}$, $\{4\ 3\}$ and $\{3\}$.

Of these 10 'consecutive subsequences', only two of them adds up to a figure that is a multiple of 4 — $\{1\ 4\ 3\}$ and $\{4\}$.

Input

The first line of input is an integer T ($T < 500$) that indicates the number of test cases. Each case consists of 5 integers A , B , C , M and N . A , B and C will be non-negative integers not greater than 1000. N and M will be a positive integers not greater than 10000.

Output

For each case, output the case number followed by the result.

Sample Input

```

2
2 1 2 4 4
923 278 195 8685 793

```

Sample Output

```

Case 1: 2
Case 2: 34

```