

11891 Deep in Jungles

The *Army of Computing Men* (ACM) is one the strongest armies in the world. They are well-equipped and have trained and armed programmers. They fight to spread creativity, teamwork, and innovation among people.

The last victory of the ACM against knowledge monopolies brought them wealthy spoils. The supply arms have located positions of N crates along a straight line in the jungle. Some trucks are to carry the crates to the army base. Since the jungle is so dense, it is costly for a truck to cut through and bring the crates back. Therefore, commanders have decided to group the crates together in points and use the minimum number of trucks. For each such gathering point, we should use a truck. Each truck has a maximum capacity of K crates and has required fuel to bring only one shipment.

The army men want to obey their commanders with minimum effort. Each crate has a weight. The effort needed to handle a crate along the line can be formulated by the weight of the crate multiplied by the distance it is carried. It is important not to change the order of crates while they are carried.

You are appointed to manage the shipping procedures. Higher ranked commanders asked you for a report on minimum needed resources. You should first consider minimizing the number of trucks and then the army effort. Write your report as soon as possible.



Input

In the first line there will be T ($T \leq 50$), number of tests. Each test begins with an integer N and K ($1 \leq N \leq 10^5$, $1 \leq K \leq 100$). It is followed by N pairs of integers p_i and w_i ($1 \leq p_i, w_i \leq 10^6$), position and the weight of i -th crate. Crates appear in the input in increasing order of their position. No two crates occupy the same position in the beginning.

Output

For each test print the minimum number of trucks and the minimum effort needed to collect all crates in a single line separated by a single space.

Hint:

In the first sample test, we need at least 3 trucks to carry the crates. The optimal solution is to carry crate number (1), (2, 3), (4, 5) in separate trucks with costs 0, 2 (for carrying crate 2 to the position of crate 3) and 8 (for carrying crate 5 to the position of crate 4) respectively.

Sample Input

```
2
5 2
1 1
4 2
```

5 10
6 10
10 2
5 5
1 1
4 2
5 10
6 10
10 2

Sample Output

3 10
1 26