

# 12000 K-Transformed Permutations

Consider a sequence of  $n$  integers  $\langle 1\ 2\ 3\ 4\ \dots\ n \rangle$ . Since all the values are distinct, we know that there are  $n$  factorial permutations. A permutation is called ***K-transformed*** if the absolute difference between the original position and the new position of every element is at most  $K$ .

Given  $n$  and  $K$ , you have to find out the total number of ***K-transformed*** permutations.

**Example:**  $n = 4, K = 2$

	1 2 3 4 (position)	Valid	Annotation
$P_1$	1 2 3 4	Yes	The original sequence. All the elements are in their original position
$P_2$	1 2 4 3	Yes	3 and 4 are reordered, but each is shifted by 1 position only.
$P_3$	1 3 2 4	Yes	
$P_4$	1 3 4 2	Yes	2 is shifted by 2 positions. $2 \leq K$ , so it's a valid one.
$P_5$	1 4 2 3	Yes	
$P_6$	1 4 3 2	Yes	
$P_7$	2 1 3 4	Yes	
$P_8$	2 1 4 3	Yes	
$P_9$	2 3 1 4	Yes	
$P_{10}$	2 3 4 1	No	1 is shifted by 3 positions. $3 > K$ and so this is an invalid permutation
$P_{11}$	2 4 1 3	Yes	
$P_{12}$	2 4 3 1	No	
$P_{13}$	3 1 2 4	Yes	
$P_{14}$	3 1 4 2	Yes	
$P_{15}$	3 2 1 4	Yes	
$P_{16}$	3 2 4 1	No	
$P_{17}$	3 4 1 2	Yes	
$P_{18}$	3 4 2 1	No	
$P_{19}$	4 1 2 3	No	4 is shifted by 3 positions. $3 > K$ and so this is also invalid
$P_{20}$	4 1 3 2	No	
$P_{21}$	4 2 1 3	No	
$P_{22}$	4 2 3 1	No	
$P_{23}$	4 3 1 2	No	
$P_{24}$	4 3 2 1	No	Here, both 4 and 1 are breaking the property.

So, for the above case, there are 14 ***2-transformed*** permutations.

## Input

The first line of input is an integer  $T$  ( $T < 20$ ) that indicates the number of test cases. Each case consists of a line containing two integers  $n$  and  $K$ . ( $1 \leq n \leq 10^9$ ) and ( $0 \leq K \leq 3$ ).

## Output

For each case, output the case number first followed by the required result. Since the result could be huge, output result modulo 73405.

**Sample Input**

```
3
4 2
100 0
10 1
```

**Sample Output**

```
Case 1: 14
Case 2: 1
Case 3: 89
```