


## 12892 Light Combat Aircraft

In graph theory, the **lowest common ancestor (LCA)** of two distinct nodes  $v$  and  $w$  in a rooted tree is the lowest (i.e. deepest) node that has both  $v$  and  $w$  as descendants, where we define each node to be a descendant of itself (so if  $v$  has a direct connection from  $w$ ,  $w$  is the lowest common ancestor).



For example, on the above tree (depicted from case 1)  $LCA(3, 5) = 1$ ,  $LCA(7, 10) = 5$ ,  $LCA(6, 5) = 5$ , etc.

In this problem, given a Forest, i.e. a disjoint union of rooted trees, you have to find out for each node  $u$  how many distinct pair of nodes  $(v, w)$  exist such that  $LCA(v, w)$  would be  $u$ . You should assume that both  $(v, w)$  and  $(w, v)$  are same pair.

### Input

First line of input file contains number of test cases,  $T \leq 100$  and  $T$  cases follow. Each case starts with an integer  $N$  ( $1 \leq N \leq 10000$ ), number of nodes in the forest. Next line contains  $N$  integers,  $p_1, p_2, \dots, p_N$  ( $0 \leq p_i \leq N$ ), where  $p_i$  is the parent of  $i$ -th ( $1 \leq i \leq N$ ) node in a rooted tree of the forest. If  $p_i = 0$  then node  $i$  is a root in rooted tree.

### Output

For each case, print the forest number starting from 1 and number of LCA pair for each node (ordered by node number) separated by space. See the sample output for exact formatting.

### Output Explanation

In **case 2**, in the given forest among the two trees rooted at 2 and 3, there is no pair for which  $LCA$  is 1 or 3. For pair  $(1, 2)$   $LCA$  is 2. So, total pair for 2 is 1.

In **case 3**, for pair  $(1, 2)$ ,  $(1, 3)$ ,  $(1, 4)$ ,  $(2, 4)$ ,  $(3, 4)$   $LCA$  is 1. For only pair  $(2, 3)$   $LCA$  is 2. There is no pair for which  $LCA$  is 3 or 4.

### Sample Input

```

4
10
  
```

```
0 1 2 1 1 5 6 6 8 5
3
2 0 0
4
0 1 2 1
4
0 1 0 3
```

### Sample Output

```
Forest#1: 29 1 0 0 9 5 0 1 0 0
Forest#2: 0 1 0
Forest#3: 5 1 0 0
Forest#4: 1 0 1 0
```